

Statistical anisotropy of magnetohydrodynamic turbulence

Wolf-Christian Müller, Dieter Biskamp

Centre for Interdisciplinary Plasma Science,

Max-Planck-Institut für Plasmaphysik, Euratom Assoziation, 85748 Garching, Germany

Roland Grappin

Observatoire de Paris-Meudon, LUTH, 92195 Meudon, France

Direct numerical simulations of decaying and forced magnetohydrodynamic (MHD) turbulence without and with mean magnetic field are analyzed by higher-order two-point statistics. The turbulence exhibits statistical anisotropy with respect to the direction of the local magnetic field even in the case of global isotropy. A mean magnetic field reduces the parallel-field dynamics while in the perpendicular direction a gradual transition towards two-dimensional MHD turbulence is observed with $k^{-3/2}$ inertial-range scaling of the perpendicular energy spectrum. An intermittency model based on the Log-Poisson approach, $\zeta_p = p/g^2 + 1 - (1/g)^{p/g}$, is able to describe the observed structure function scalings.

PACS: 47.27Gs; 47.65+a; 47.27Eq; 52.35.Ra

Turbulence is the natural state of many plasma flows observed throughout the universe, its statistical properties being essential for the theoretical understanding of, e.g., star-forming regions in the interstellar medium, the convection in planetary and stellar interiors, as well as the dynamics of stellar winds. The solar wind, in particular, represents the only source of in-situ measurements, since laboratory experiments are far from generating fully-developed turbulence at high magnetic Reynolds numbers. A simplified nonlinear model of turbulent plasmas is incompressible magnetohydrodynamics (MHD) [1]. In this framework the kinetic nature of microscopic processes responsible for, e.g., energy dissipation, is neglected when studying the fluid-like macroscopic plasma motions.

The spatial similarity of incompressible MHD turbulence is usually investigated by considering two-point statistics of the Elsässer variables $\mathbf{z}^\pm = \mathbf{v} \pm \mathbf{B}$ [2] combining velocity $\mathbf{v}$ and magnetic field $\mathbf{B}$ (given in Alfvén-speed units). Restricting consideration to turbulence with small cross helicity $H^C = \int_V dV (\mathbf{v} \cdot \mathbf{B})$, V being the volume of the system, allows to set $\mathbf{z}^+ \simeq \mathbf{z}^- = \mathbf{z}$. With $\delta z_\ell = [\mathbf{z}(\mathbf{r} + \boldsymbol{\ell}) - \mathbf{z}(\mathbf{r})] \cdot \boldsymbol{\ell}/\ell$ the longitudinal isotropic structure functions of order p are defined as $S_p(\ell) = \langle \delta z_\ell^p \rangle$, the angular brackets denoting spatial averaging. The structure functions exhibit self-similar scaling $S_p(\ell) \sim \ell^{\zeta_p}$ in the inertial range where the dynamical influence of dissipation, turbulence driving and system boundaries is weak.

The inertial range has been introduced in Kolmogorov's K41 phenomenology of incompressible hy-

drodynamic turbulence [3,4] which assumes a spectral energy-cascade driven by the break-up of turbulent eddies. This leads to the experimentally well-verified energy-spectrum $E(k) \sim k^{-5/3}$ [5] corresponding to $\zeta_2 = 2/3$. Iroshnikov and Kraichnan (IK) [6,7] included the effect of a magnetic field by founding the energy-cascade on the mutual scattering of Alfvén waves triggered by velocity fluctuations. The IK picture phenomenologically yields $E(k) \sim k^{-3/2}$, i.e., $\zeta_2 = 1/2$.

The validity of the two phenomenologies in MHD turbulence is still under discussion. Two-dimensional direct numerical simulations (DNS) support the IK picture [8,9] while three-dimensional simulations exhibit K41 scaling behavior [10]. Analytical results [11] also suggest $\zeta_3 = 1$, consistent with K41 energy-spectra measured in the solar wind [12]. A recent phenomenology of Goldreich and Sridhar [13] postulates a balance between K41 and IK energy cascades and accounts for the local anisotropy induced by $\mathbf{B}$. However, DNS which claim to support this picture [14,15] suffer from moderate numerical resolution, making the identification of self-similar scaling ranges difficult.

In this Letter we examine three-dimensional pseudospectral DNS of decaying isotropic MHD turbulence and of driven turbulent systems with mean magnetic field $\mathbf{B}_0$ at comparably high resolutions of up to 512^3 collocation points. The structure functions are found to be anisotropic with respect to the local magnetic field. The effect increases with magnetic-field strength, reducing the spatial intermittency of the turbulence in the parallel-field direction while rendering the system quasi-two-dimensional perpendicular to $\mathbf{B}$. An intermittency model based on the Log-Poisson approach agrees well with the observed structure-function scalings.

The simulations are performed by numerically solving the incompressible MHD equations

$$\partial_t \boldsymbol{\omega} - \nabla \times (\mathbf{v} \times \boldsymbol{\omega} + \mathbf{j} \times \mathbf{B}) = \mu_\nu (-1)^{\nu-1} \Delta^\nu \boldsymbol{\omega}, \quad (1)$$

$$\partial_t \mathbf{B} - \nabla \times (\mathbf{v} \times \mathbf{B}) = \eta_\nu (-1)^{\nu-1} \Delta^\nu \mathbf{B}, \quad (2)$$

$$\boldsymbol{\omega} = \nabla \times \mathbf{v}, \quad \mathbf{j} = \nabla \times \mathbf{B}, \quad \nabla \cdot \mathbf{v} = \nabla \cdot \mathbf{B} = 0,$$

in a periodic cube with a pseudo-spectral method using spherical mode-truncation to reduce aliasing effects [16]. All simulations comprise about 9 eddy-turnover times,

a turnover time being defined as the period required to reach maximal energy dissipation when starting from smooth fields. The initial conditions in the decaying case are characterized by global equipartition of kinetic and magnetic energy $E^K = E^M = 0.5$ with a spectral distribution $\sim \exp(-k^2/k_0^2)$, $k_0 = 4$, and random phases. The driven runs start from a quasi-stationary state with $E^K \simeq 0.75$, $E^M \simeq 0.8$ generated by forcing the system for 24 turnover times. The forcing is realized by freezing all modes in a sphere of radius $k_f = 2$, allowing energy transfer to higher wavenumbers by nonlinear interactions only. The magnetic helicity in all simulations is finite, $H^M = \int_V dV (\mathbf{A} \cdot \mathbf{B}) \approx 0.7 H_{\max}^M$ with the magnetic vector potential $\mathbf{A}$ and $H_{\max}^M \sim E_{\max}^M/k_0$. For the driven cases the $\mathbf{B}_0$ -component which renders H^M gauge-dependent has been subtracted and $k_0 \simeq k_f$. The cross helicity H^C is approximately zero in the decaying system and $\simeq 0.12$ for the driven runs, mildly fluctuating around the respective value. The generalized magnetic Prandtl number μ_ν/η_ν is equal to unity with hyperdiffusive dissipation operators $\nu = 2$, $\mu_2 = \eta_2 = 3 \times 10^{-8}$. Test runs with $\nu = 1$ show no notable difference in the reported results.

Weak-turbulence theory [17] and numerical simulations [18–20] show that the IK cascade is spatially anisotropic, the underlying three-wave interactions distributing energy predominantly perpendicular to $\mathbf{B}$. The associated spectral dependence on a mean magnetic field has been studied numerically as well as in the reduced-MHD framework ([21] and references therein) while statistical anisotropy has been found in the second-order structure functions of MHD turbulence [22]. In order to assess this anisotropy by higher-order statistics, parallel and perpendicular structure functions are calculated with field increments, δz_ℓ , taken in the corresponding directions with respect to the local magnetic field. This is in contrast to isotropic structure functions where the δz_ℓ are measured without preferred direction. The local magnetic field on the increment length-scale, defining field-parallel and field-perpendicular directions at each spatial co-ordinate, is found by applying a top-hat filter of width ℓ , i.e., by multiplying $\mathbf{B}$ with the Fourier-space filter kernel $G_\ell(k) = \sin(k\ell/2)/(k\ell/2)$. The obtained structure functions, computed using $|\delta z_\ell|$ to avoid cancelation effects, exhibit inertial-range scaling with exponents ζ_p which can be determined up to order $p = 8$ by a saddle-point in the logarithmic derivative. The results have been cross-checked using the extended self-similarity property [23] of the data.

Fig. 1 shows that the decaying system with $B_0 = 0$ (512^3 collocation points) is statistically anisotropic. The field-perpendicular scalings display increased intermittency compared to the isotropic structure functions. Assuming that the formation of intermittent structures primarily depends on the turbulent energy available at the corresponding spatial scales, the observation is consistent with both the IK and the K41 cascade being strongest

perpendicular to $\mathbf{B}$. The former due to the inherent anisotropy of Alfvén-wave scattering, the latter because field lines resist bending but can be shuffled easily by eddies perpendicular to the local field. Consequently, the field-parallel energy transfer is depleted leading to less intermittent parallel scalings.

The dependence of statistical anisotropy on the magnetic-field strength is examined in forced-turbulence simulations with mean magnetic field ($B_0 = 5, 10$) allowing a reduction of numerical resolution in the mean-field direction to respectively 256 and 128 collocation points (cf. Fig. 2). The quasi-stationary forced systems assume constant energy ratios of mean field to turbulent fluctuations. The parallel scalings shown in Fig. 1 display decreasing intermittency with raising B_0 , i.e., an asymptotic approach towards a straight line. Referring to Kolmogorov’s refined-similarity hypotheses [24], this is equivalent to higher parallel-field homogeneity of small-scale dissipative structures, i.e., current and vorticity micro-sheets, due to their stronger alignment along $\mathbf{B}_0$. The perpendicular statistics (Fig. 1) become increasingly two-dimensional, getting close to values found in DNS of two-dimensional MHD turbulence (The exponents for $B_0 = 10$ and those obtained from 2D-DNS [9] coincide within the error-margin.). The asymptotic state results from micro-sheet alignment along $\mathbf{B}_0$ which decreases the spatial extent of the sheets in the field-perpendicular direction to quasi-one-dimensional dissipative ribbons.

The ζ_2 -exponents are related to the inertial-range scaling of the one-dimensional energy spectra $E_k = 1/2 \int dk_1 \int dk_2 (|\mathbf{v}_k|^2 + |\mathbf{B}_k|^2)$ with k_1, k_2 spanning planes perpendicular to the component of $\mathbf{k}$ associated with the spatial increment ℓ . For the field-parallel and field-perpendicular spectra this gives $E_k^{\parallel, \perp} \sim k^{-(1+\zeta_2^{\parallel, \perp})}$. Fig. 1 yields $E_k^{\parallel}$ -exponents in the range $[-1.8, -1.9]$ while the $E_k^{\perp}$ -scaling changes from K41, $\sim k^{-5/3}$, to IK, $\sim k^{-3/2}$, with increasing B_0 . This agrees with DNS of two-dimensional MHD turbulence and suggests that, contrary to the three-dimensional case where K41 scaling is observed, the restriction to a quasi-two-dimensional geometry increases the importance of the inherently two-dimensional Alfvén-wave cascade (IK) compared to the eddy-breakup process (K41).

In Fig. 2, $E_k^{\parallel}$ and $E_k^{\perp}$ with respect to the fixed $\mathbf{B}_0$ -axis are given for $B_0 = 0, 5, 10$. The spectra are time-averaged over 4 eddy turnover-times and normalized in amplitude assuming a K41 dependence on the mean energy-dissipation $\varepsilon = -\dot{E}$ as $\sim \varepsilon^{2/3}$. Wavenumbers are normalized with the generalized K41 dissipation-length $\ell_D = (\mu^3/\varepsilon)^{1/(6\nu-2)}$. The normalization, though unnecessary for the driven runs, allows comparison with the decaying case shown in the inset in Fig. 2. For $B_0 = 0$ the parallel and perpendicular energy spectra differ slightly at largest scales where the few involved Fourier modes do not isotropize perfectly. The inertial range exhibits

K41 scaling which leads to a clear deviation from the horizontal under the applied $k^{3/2}$ compensation. For finite B_0 the spectra display a marked anisotropy in agreement with the perpendicular and parallel structure functions. With growing B_0 the $E_k^\perp$ asymptotically follow IK scaling while the $E_k^\parallel$ indicate an increasing depletion of small-scale turbulence. The field-parallel damping results from the stiffness of magnetic field lines in agreement with the picture of field-aligned dissipative structures. This corresponds to an increase of the associated dissipation length [25]. The different amplitudes of $E_k^\perp$ and $E_k^\parallel$ beyond the forcing range $k \gtrsim 0.02$ have been found similarly in shell-model calculations of anisotropic MHD turbulence [26] resulting from an equilibrium between field-perpendicular and isotropic energy cascades.

Intermittency, the departure of turbulence from strict spatial self-similarity, leads to ‘anomalous’ non-linear behavior of the ζ_p . The Log-Poisson intermittency model [27] reproduces these experimental and numerical findings in hydrodynamic and MHD turbulence very well. Its generic form $\zeta_p = (1-x)p/g + C_0(1 - [1-x/C_0]^{p/g})$ [28,29] depends on the codimension C_0 of the most singular dissipative structures (in three-dimensions $C_0 = 2$ for filaments, $C_0 = 1$ for micro-sheets), the scale dependence of dissipation in these structures $\varepsilon_\ell^{(\infty)} \sim \ell^{-x}$ and the phenomenological non-intermittent scaling $\delta z_\ell \sim \ell^{1/g}$ ($g = 3$ for K41, $g = 4$ for IK). Usually, x and g are linked by assuming equal scaling of the timescale t_ℓ^∞ of $\varepsilon_\ell^{(\infty)} \sim E^\infty/t_\ell^\infty$ and the nonlinear transfer time t_ℓ^{NL} of the energy cascade $\varepsilon \sim \delta z_\ell^2/t_\ell^{\text{NL}}$ yielding $x = 2/g$. Here, E^∞ denotes the amount of energy dissipated in the most singular structures. The successful hydrodynamic She-Lévêque formula [27] results from $C_0 = 2$, $g = 3$ while isotropic structure-function scalings in DNS of three-dimensional MHD turbulence are well reproduced with $C_0 = 1$, $g = 3$ [30].

To model statistical anisotropy we extend the approach presented in [30] by dropping the plausible but not mandatory scaling-equality of t_ℓ^{NL} and t_ℓ^∞ . Instead, t_ℓ^∞ is fixed to the K41 timescale, $t_\ell^\infty \sim \ell/\delta z_\ell \sim \ell^{1-1/g}$, which together with $C_0 = 1$ leads to

$$\zeta_p = p/g^2 + 1 - (1/g)^{p/g}. \quad (3)$$

Fig. 3 shows the predictions of Eq. (3) with the corresponding numerical values of g . The isotropic MHD intermittency model based on K41 scaling [30] is denoted by the solid line in Figs. 1 and 3. For increasing B_0 the limiting value of g in the parallel direction is 1, standing for spatially homogeneous dissipation. The asymptotic perpendicular exponents should be reproduced by $g = 4$ to be consistent with the IK scaling observed in this work and in DNS of two-dimensional MHD turbulence. The fact that the observed perpendicular exponents for $B_0 = 10$ correspond to the model with $g \approx 4.4$ can be

ascribed to the simplicity of the approach which nevertheless captures the basic physics of the system.

By detaching t_ℓ^{NL} and t_ℓ^∞ the strengths of field-perpendicular and parallel cascades can be modified without affecting the mechanism of most singular dissipation. The quantity $g/3$ expresses the cascade strength relative to the isotropic K41 case as can be seen by writing a modified K41 transfer-time $t_\ell^{\text{NL}} \sim (\ell/\ell_0)^x (\ell/\delta z_\ell)$ introducing an arbitrary reference length ℓ_0 and the dimensionless efficiency parameter χ . Combination with $\delta z_\ell^2/t_\ell^{\text{NL}} = \text{const.}$ yields $t_\ell^{\text{NL}} \sim \ell^{(1+\chi)2/3}$ compared to the standard-phenomenology result $\sim \ell^{2/3}$. A value of $\chi = 0$ ($g = 3$) yields the isotropic K41 case while $\chi > 0$ ($g < 3$) corresponds to cascade depletion and $\chi < 0$ ($g > 3$) to cascade enhancement. In this picture the cascade efficiency is controlled by the factor $(\ell/\ell_0)^x$ in t_ℓ^{NL} , lumping together deviations of the physical transfer-process from the K41 picture *and* differences in the amount of cascading energy compared to the isotropic case. For example, the model indicates a growing field-perpendicular cascade with increasing B_0 though scalings suggest a transition from K41 to the less efficient IK cascade mechanism. This efficiency reduction is, however, over-compensated by the increase of energy cascading field-perpendicularly compared to the isotropic situation. The model reproduces the numerical data very well and in agreement with the physical interpretation suggested above. With increasing B_0 a larger fraction of energy compared to the isotropic case ($B_0 = 0$) is spectrally transferred perpendicular to the magnetic field while the cascade becomes successively damped in the parallel-field direction.

In summary, we have analyzed DNS of decaying and forced MHD turbulence without and with varying mean magnetic field using higher-order statistics. Globally isotropic turbulence exhibits statistical anisotropy, attributed to the influence of the local magnetic field on the nonlinear energy cascade. An increasing mean magnetic-field B_0 damps the parallel-field dynamics while in the perpendicular direction a gradual transition towards two-dimensional MHD turbulence is observed with perpendicular energy-spectra showing IK scaling. A modified Log-Poisson intermittency model reproduces the statistical anisotropy by phenomenological tuning of the respective energy cascades. The anisotropic approach of Goldreich and Sridhar, therefore, seems to be a promising concept though the proposed realizations for weak, ‘intermediate’ and ‘strong’ turbulence remain questionable.

-
- [1] D. Biskamp, *Nonlinear Magnetohydrodynamics* (Cambridge University Press, Cambridge, 1993).
 - [2] W. M. Elsässer, *Physical Review* **79**, 183 (1950).

- [3] A. N. Kolmogorov, Proceedings of the Royal Society A **434**, 9 (1991), [Dokl. Akad. Nauk SSSR, 30(4), 1941].
- [4] A. N. Kolmogorov, Proceedings of the Royal Society A **434**, 15 (1991), [Dokl. Akad. Nauk SSSR, 32(1), 1941].
- [5] U. Frisch, *Turbulence* (Cambridge University Press, Cambridge, 1996).
- [6] P. S. Iroshnikov, Soviet Astronomy **7**, 566 (1964), [Astron. Zh., 40:742, 1963].
- [7] R. H. Kraichnan, Physics of Fluids **8**, 1385 (1965).
- [8] D. Biskamp and E. Schwarz, Physics of Plasmas **8**, 3282 (2001).
- [9] H. Politano, A. Pouquet, and V. Carbone, Europhysics Letters **43**, 516 (1998).
- [10] D. Biskamp and W.-C. Müller, Physics of Plasmas **7**, 4889 (2000).
- [11] H. Politano and A. Pouquet, Geophysical Research Letters **25**, 273 (1998).
- [12] R. J. Leamon *et al.*, Journal of Geophysical Research **103**, 4775 (1998).
- [13] P. Goldreich and S. Sridhar, Astrophysical Journal **438**, 763 (1995).
- [14] J. Maron and P. Goldreich, Astrophysical Journal **554**, 1175 (2001).
- [15] J. Cho, A. Lazarian, and E. T. Vishniac, Astrophysical Journal **564**, 291 (2002).
- [16] A. Vincent and M. Meneguzzi, Journal of Fluid Mechanics **225**, 1 (1991).
- [17] S. Galtier, S. V. Nazarenko, A. C. Newell, and A. Pouquet, Journal of Plasma Physics **63**, 447 (2000).
- [18] J. V. Shebalin, W. H. Matthaeus, and D. Montgomery, Journal of Plasma Physics **29**, 525 (1983).
- [19] R. Grappin, Physics of Fluids **29**, 2433 (1986).
- [20] C. S. Ng and A. Bhattacharjee, Astrophysical Journal **465**, 845 (1996).
- [21] S. Oughton, W. H. Matthaeus, and S. Ghosh, Physics of Plasmas **5**, 4235 (1998).
- [22] L. J. Milano, W. H. Matthaeus, P. Dmitruk, and D. C. Montgomery, Physics of Plasmas **8**, 2673 (2001).
- [23] R. Benzi *et al.*, Physical Review E **48**, R29 (1993).
- [24] A. N. Kolmogorov, Journal of Fluid Mechanics **13**, 82 (1962).
- [25] R. Grappin and A. Mangeney, in *Proceedings of the Eighth International Solar Wind Conference* (AIP Conference Proceedings, Dana Point, CA, 1996), Vol. 382, pp. 250–255.
- [26] V. Carbone and P. Veltri, Geophysical and Astrophysical Fluid Dynamics **52**, 153 (1990).
- [27] Z.-S. She and E. Lévêque, Physical Review Letters **72**, 336 (1994).
- [28] R. Grauer, J. Krug, and C. Marliani, Physics Letters A **195**, 335 (1994).
- [29] H. Politano and A. Pouquet, Physical Review E **52**, 636 (1995).
- [30] W.-C. Müller and D. Biskamp, Physical Review Letters **84**, 475 (2000).

List of Figures

- 1 Scaling exponents ζ_p of perpendicular (filled symbols) and parallel (open symbols) structure-functions $S_p(\ell) = \langle |\delta z_\ell|^p \rangle$ for $B_0 = 0, 5, 10$ (resp. circles, diamonds, triangles) together with isotropic scalings from 3D-DNS (solid line, [30]). Error-bars are given in Fig. 3. 4
- 2 Normalized, time-averaged parallel (dashed) and perpendicular (solid) energy spectra compensated with $k^{3/2}$ for $B_0 = 0$ (inset), $B_0 = 5$ (lower solid line, upper dashed line) and $B_0 = 10$ 4
- 3 Scaling results as in Fig. 1 combined with predictions of Eq. (3) (dotted lines). The numerical values of g are given next to the respective curves, $g = 3$ corresponds to the isotropic MHD intermittency model [30]. . 5




FIG. 1.




FIG. 2.




FIG. 3.